

T7 RNA Polymerase

Catalog Number LDG0001RI

Package

er LDG0001RI

25,000 U / Customized package

For full product information, images and publications, please visit our website.



Overview

Description

Bacteriophage T7 RNA Polymerase is a DNA-dependent RNA polymerase with high specificity for the T7 promoter. This enzyme catalyzes the $5' \rightarrow 3'$ synthesis of RNA from DNA downstream from its promoter.

Components

Package	ltems	Quantity
	T7 RNA Polymerase (200 U/μL)	1 vial (25,000 U)
25,000 U	10× RNA Polymerase reaction buffer	1 vial (1 mL)
	100 mM DTT	1 vial (1 mL)

Specifications

Expression System	Storage Buffer
Escherichia coli	T7 RNA Polymerase is supplied in 100 mM Tris- HCl (pH 7.9), 20 mM KCl, 1 mM DTT, 1 mM EDTA, 0.1% Triton® X-100 and 50% (v/v) glycerol.
Purity >98% as determined by SDS-PAGE analysis.	Unit Definition One unit is defined as the amount of the enzyme incorporates 1 nmol of ATP into acid-insoluble product in 1 hour at 37°C.

Tainan Headquarter

Taipei Office

ⓒ +886-6-2536677 ⓓ +886-6-2531536

☑ info@leadgene.com.tw

& +886-2-27065528



Reaction Condition

1× RNA Polymerase Reaction Buffer, supplemented with 0.5 mM each ATP, UTP, GTP, CTP, and DNA template containing the T7 RNA Polymerase promoter. Incubate at 37°C. 10× RNA Polymerase Reaction Buffer: 400 mM Tris-HCl (pH 8.0), 60 mM MgCl₂, and 20 mM spermidine.

Instruction

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage

This product is stable after storage at:

- -20°C or -80°C for 12 months under sterile conditions from date of receipt.
- For optimal storage, aliquot the enzyme, reaction buffer and DTT reagent into smaller quantities and store at recommended temperature.
- For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarter

Taipei Office

☑ info@leadgene.com.tw

& +886-2-27065528