

Anti-DENV M Antibody [Clone 31-1]

Catalog Number	LDG0009YA
Package	100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Overview

Description

Dengue virus belongs to the family Flaviviridae, genus Flavivirus and consists of four distinct serotypes (DENV1 to 4). Dengue virus is transmitted by mosquitos majorly of species Aedes aegypti and Aedes Albopictus that widespread around tropics and subtropics. The Membrane protein is the second smallest protein of dengue virus. Before dengue virus particles release, host protease furin cleaves prM into M. This amphipathic helical portion acts as an important component of the mature virion, interacting with envelope protein to promote invasion, packing mature, and release.

Product Note

Recommended dilution factor:

ELISA: 1:5000-20000

WB: 1:1000-5000

IFA: 1:500-1000

FACS: Assay dependent

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

Specifications

Host

Mouse

Clonality

Monoclonal

Isotype

IgG2b

Clone Name

clone 31-1

Immunogen

Dengue virus M

Reactivity

Dengue virus

Tainan Headquarters

+886-6-2536677

bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677

Application

ELISA, WB, IFA, FACS

Concentration

1 mg/mL

Specificity

M (membrane) protein

Conjugation

Unconjugated

Storage Buffer

Phosphate Buffered Saline containing 0.03% ProClin 300, pH 7.4.

Form

Liquid

Instruction**Shipping**

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.



Stability & Storage


This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.


Avoid repeated freeze/thaw cycles.

Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.


Image**Tainan Headquarters** +886-6-2536677 bd@leadgene.com.tw**Innovation & Research Center** +886-2-27065528**CLD Center** +886-6-2536677




Immunofluorescence analysis of Anti-DENV M Antibody [Clone 31-1]
 Control Vero cells (uninfected) and Virus (DENV 1,2) infection Vero cells were fixed in 4% PFA, permeabilized with PBS containing 0.1% Triton X-100. Cells were stained with mouse anti-DENV M monoclonal antibody (1:400) followed by secondary antibodies (goat anti-Mouse IgG-iFluor 488, 1:400, green) and cell nuclei were stained with Hoechst 33342 (Blue).



Immunofluorescence analysis of Anti-DENV M Antibody [Clone 31-1]
 Control BHK cells (uninfected) and Virus (DENV 1,2) infection BHK cells were fixed in 4% PFA, permeabilized with PBS containing 0.1% Triton X-100. Cells were stained with mouse anti-DENV M monoclonal antibody (1:400) followed by secondary antibodies (goat anti-Mouse IgG-iFluor 488, 1:400, green) and cell nuclei were stained with Hoechst 33342 (Blue).



ELISA titration of Anti-DENV M Antibody [Clone 31-1]
 Titration curve of anti-DENV M antibody in ELISA. Red: DENV M; Black: BSA (negative control).



Western blotting analysis of Anti-DENV M Antibody [Clone 31-1]
 Virus infection C636 cell lysates were stained with mouse anti-DENV M monoclonal antibody at 1:1000 dilution.

Lane 1: ZIKV infection (10 µg), Lane 2: DENV1 infection (10 µg), Lane 3: DENV2 infection (10 µg), Lane 4: DENV3 infection (10 µg), Lane 5: DENV4 infection (10 µg), Lane 6: C636 cell lysate (10 µg).



Western blotting analysis of Anti-DENV M Antibody [Clone 31-1]
 Virus infection Vero cell lysates were stained with mouse anti-DENV M monoclonal antibody at 1:1000 dilution.

Lane 1: Vero cell lysate (10 µg), Lane 2: ZIKV infection (10 µg), Lane 3: DENV1 infection (10 µg), Lane 4: DENV2 infection (10 µg), Lane 5: DENV3 infection (10 µg), Lane 6: DENV4 infection (10 µg).

Disclaimer : For Research Use or Further Manufacturing Only.