

## AccuBlot<sup>™</sup> Anti-Rabbit IgG (H+L) Antibody (HRP)

Catalog Number	LDG0015YE
Package	100 μL / 1 mL / Customized package

For full product information, images and publications, please visit our website.



### **Overview**

#### Description

Leadgene AccuBlotTM Anti-Rabbit IgG (H+L) Antibody (HRP) is conjugated with high-activity HRP. In coimmunoprecipitation (co-IP) assay, denatured IgG from IP samples may show interference and result in immunoblotting at ~55 kDa (heavy chain) and ~25 kDa (light chain) by using traditional secondary antibodies. The products of AccuBlotTM series are suitable for native IgG detection.

#### **Product Note**

Recommended dilution factor:

ELISA: 1:2500-10000 WB: 1:1000-3000

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

# **Specifications**

Host Clonality Monoclonal Mouse Immunogen Isotype lgG2b Purified rabbit IgG Reactivity **Application** ELISA, WB Native rabbit IgG Storage Buffer Conjugation HRP Phosphate Buffered Saline containing 1% BSA, 0.03% Proclin300 and 50% glycerol, pH 7.4.

**Tainan Headquarters** 

**Innovation & Research Center** 

**CLD Center** 



Specificity Form  IgG Liquid		
	Specificity	Form
		Liquid

### Instruction

### Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

### Stability & Storage

This product is stable after storage at:


- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles and protect from light.

Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.

### **Image**





Western blot analysis of native rabbit IgG (lane 1) and denatured rabbit IgG (lane 2), using AccuBlot™ anti-Rabbit IgG-HRP at a 1:5000 dilution.

**Disclaimer:** For Research Use or Further Manufacturing Only.