

Goat Anti-Human IgG (Fc Fragment) Antibody (HRP)

Catalog Number	LDG0017YE
Package	1 mL / Customized package

For full product information, images and publications, please visit [our website](#).



Overview

Description

Leadgene Goat anti-Human IgG, Fc Fragment is purified from goat antiserum by immunoaffinity chromatography and conjugated with high-activity HRP. Minimal cross-reactivity with other isotypes or species.

Product Note

Recommended dilution factor:

ELISA: 1:5000-25000

IHC: 1:500-2500

WB: 1:5000-25000

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

Specifications

Host

Goat

Clonality

Polyclonal

Immunogen

Gamma immunoglobulin Fc region

Reactivity

Human

Application

ELISA, IHC, WB

Conjugation

HRP

Tainan Headquarters

+886-6-2536677

bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677

Storage Buffer

Phosphate Buffered Saline containing 1% BSA, 0.03% ProClin 300 and 10% glycerol, pH 7.4.

Form

Liquid

Specificity

IgG, Fc Fragment

Instruction**Shipping**

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage

This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles and protect from light.

Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarters

+886-6-2536677

bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677