

Lactate Oxidase (LOX)

Catalog Number	LDG0033RG
Package	Customized package

For full product information, images and publications, please visit our website.



Specifications

Expression System	Activity
Escherichia coli	≥200 U/mg
Unit Definition One unit causes the formation of one micromole of hydrogen peroxide (half a micromole of quinoneimine dye) per minute under the conditions described below.	Form Lyophilized (Yellowish amorphous powder)

Background

Synonyms

LOX, LctO, Lactic oxygenase, Lactic oxidase, Lactate monooxygenase, L-lactate oxidase

Instruction

Reconstitution

It is recommended to weight and reconstitute 10 mg of lyophilized powder in 200 μ L double-distilled water directly (final activity is 10 U/ μ L) and incubate the solution for at least 10 mins to ensure sufficient redissolved.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Tainan Headquarters

Innovation & Research Center

CLD Center

& +886-6-2536677

® +886-2-27065528

& +886-6-2<u>5</u>36677

☑ bd@leadgene.com.tw



Datasheet

Stability & Storage

This product is stable at -20°C for long-term storage under sterile conditions.

Avoid repeated free-thaw cycles.

Image







pH stability of LOX.

The enzyme powder was reconstituted by double-distilled water and treated with different pH buffer condition at 25°C for 16 hours. pH 4.0-6.0, 0.1 M Sodium citrate buffer; pH 7.5-8.0, 0.1 M Potassium phosphate buffer; pH 8.5, 0.1 M Tris-HCI buffer; pH 10.0-11.0, 0.1 M Carbonate-bicarbonate buffer. Thermal stability of LOX.

The enzyme powder was reconstituted by double-distilled water and treated with different temperatures for 10 minutes. Final concentration: 10 U/ mL. pH activity of LOX.

The buffer conditions with various pH values were used in the reaction at 37°C. pH 4.0-6.0, 0.1 M Sodium citrate buffer; pH 7.5-8.0, 0.1 M Potassium phosphate buffer; pH 8.5, 0.1 M Tris-HCl buffer; pH 10.0-11.0, 0.1 M Carbonate-bicarbonate buffer.

Tainan Headquarters

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-6-2536677

& +886-2-27065528

☑ bd@leadgene.com.tw





Temperature activity of LOX. The enzyme reactions in 20 M K-Phosphate buffer, pH 7.5, were carried out under different temperatures.

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarters

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw