

## Human IL-8 (aa 28-99), His Tag, E. coli

<b>Catalog Number</b>	LDG010PHE
<b>Package</b>	5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



### Specifications

#### Species of Origin

Human

#### Affinity Tag

His Tag (C-term)

#### Purity

>98% as determined by SDS-PAGE analysis.

#### Activity

Measure by its ability to chemoattract PBMC. The ED<sub>50</sub> for this effect is <2 ng/mL.

#### Form

Lyophilized

#### Expression System

Escherichia coli

#### Storage Buffer

Lyophilized from a 0.2 µm filtered solution of PBS, pH 8.0.

#### Molecular weight

The protein has a calculated MW of 9.32 kDa. The protein migrates as 11 kDa under reducing condition (SDS-PAGE analysis).

#### Endotoxin Level

<0.1 EU per 1 µg of the protein by the LAL method.

### Background

#### Tainan Headquarters

+886-6-2536677

bd@leadgene.com.tw

#### Innovation & Research Center

+886-2-27065528

#### CLD Center

+886-6-2536677

### Background

Human IL-8 (aa 28-99) predicts a molecular mass of 8.5 kDa and is the 72 amino acid residue form Ser 28 to Ser 99 of mature human IL-8. This recombinant protein is the major form secreted by macrophages. IL-8 is a chemokine produced by macrophages and other cell types such as epithelial cells, airway smooth muscle cells and endothelial cells.

### Uniprot ID

#P10145

### Synonyms

C-X-C motif chemokine 8, Chemokine, C-X-C motif ligand 8, Emotakin, Granulocyte chemotactic protein 1, GCP-1, Monocyte-derived neutrophil chemotactic factor, MDNCF, Monocyte-derived neutrophil-activating peptide, MONAP, Neutrophil-activating protein 1, NAP-1, Protein 3-10C, T-cell chemotactic factor

### Sequence Note

Ser28-Ser99

## Instruction

### Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

### Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

### Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

## Image

### Tainan Headquarters

+886-6-2536677


bd@leadgene.com.tw

### Innovation & Research Center

+886-2-27065528


### CLD Center

+886-6-2536677



Measure by its ability to chemoattract PBMC using a concentration range of 1-10 ng/mL.

Activity analysis of recombinant Human IL-8 (aa 28-99), His Tag, E. coli.



SDS-PAGE analysis of recombinant human IL-8 (aa 28-99).

**Disclaimer :** For Research Use or Further Manufacturing Only.