

Human IL-17A, His Tag, E. coli

Catalog Number LDG019PHE

Package 5 μg / 20 μg / 100 μg / Customized package

For full product information, images and publications, please visit our website.



Specifications

Species of Origin

Human

Affinity Tag

His Tag (C-term)

Purity

>98% as determined by SDS-PAGE analysis.

Activity

Measure by its ability to induce IL-6 secretion in 3T3 cells. The ED $_{50}$ for this effect is <6 ng/mL.

Form

Lyophilized

Expression System

Escherichia coli

Storage Buffer

Lyophilized from a 0.2 μm filtered solution of PBS, pH 8.0.

Molecular weight

The protein has a calculated MW of 16.47 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).

Endotoxin Level

<0.01 EU per 1 μg of the protein by the LAL method.

Background



Background

Interleukin 17A (IL-17A) predicts a molecular mass of 35 kDa, encoded IL17A gene is a proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappa B and mitogenactivated protein kinases.

Uniprot ID

#Q16552

Synonyms

Cytotoxic T-lymphocyte-associated antigen 8, CTLA-8

Sequence Note

Ile20-Ala155

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H_2O to a concentration of 200 $\mu g/mL$ and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.


Avoid repeated freeze/thaw cycles.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Image





SDS-PAGE analysis of recombinant human IL-17A.

Disclaimer : For Research Use or Further Manufacturing Only.