

Mouse EpCAM, His Tag, CHO

Catalog Number	LDG020PMM
Package	5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Specifications

Species of Origin

Mouse

Expression System

CHO

Affinity Tag

His Tag (C-term)

Storage Buffer

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Purity

>95% as determined by SDS-PAGE analysis.

Molecular weight

The protein has a calculated MW of 28.46 kDa.
The protein migrates as 30-35 kDa under reducing condition (SDS-PAGE analysis).

Activity

Measured by its ability to support the adhesion of NIH-3T3 mouse embryonic fibroblast cells, with the ED50 ranging from 1 to 5 µg/mL.

Endotoxin Level

<0.1 EU per 1 µg of the protein by the LAL method.

Form

Lyophilized

Background

Tainan Headquarters

+886-6-2536677

bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677

Background

EpCAM (Epithelial Cell Adhesion Molecule), a transmembrane glycoprotein, is expressed in epithelial tissues and various carcinomas. It mediates cell-cell adhesion and signaling, influencing cell proliferation, migration, and differentiation. EpCAM is involved in maintaining epithelial integrity and regulating epithelial-mesenchymal transition (EMT) during development and tissue repair. Additionally, it serves as a diagnostic and prognostic marker in cancer, where its overexpression correlates with tumor aggressiveness and poor clinical outcomes. EpCAM-targeted therapies, including monoclonal antibodies and immunotherapies, hold promise in cancer treatment. Understanding EpCAM's roles in physiology and pathology is crucial for developing effective therapeutic strategies.

Uniprot ID

AAH05618.1

Synonyms

Epithelial cell adhesion molecule, Epithelial glycoprotein 314 (EGP314; mEGP314), Protein 289A, Tumor-associated calcium signal transducer 1, CD326

Sequence Note

Met1-Thr266

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration of 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage


This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.


Avoid repeated freeze/thaw cycles.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Image


Mouse EpCAM, His Tag, CHO (LDG020PMM) supports the adhesion of NIH-3T3 mouse embryonic fibroblast cells, with the ED50 ranging from 1 to 5 μ g/mL.



SDS-PAGE analysis of recombinant mouse EpCAM.

Disclaimer : For Research Use or Further Manufacturing Only.