

Swine FGF-2, His Tag, E. coli

Catalog Number	LDG021PSE
Package	5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Specifications

Species of Origin

Swine

Affinity Tag

His Tag (N-term)

Purity

>98% as determined by SDS-PAGE analysis.

Activity

Measure by its ability to induce proliferation in 3T3 cells. The ED₅₀ for this effect is <2 ng/mL.

Form

Lyophilized

Expression System

Escherichia coli

Storage Buffer

Lyophilized from a 0.2 µm filtered solution of PBS containing 0.01% sarkosyl, pH 7.4.

Molecular weight

The protein has a calculated MW of 18.1 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).

Endotoxin Level

<0.1 EU per 1 µg of the protein by the LAL method.

Background

Tainan Headquarters

+886-6-2536677

bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677

Background

Basic fibroblast Growth Factors (FGF-2, bFGF), a pleiotropic cytokine, plays multiple roles in different cells and tissues. FGF-2 can stimulate smooth muscle cell growth, wound healing, and tissue repair. In addition, FGF-2 has been shown to regulate the generation of neurons and astrocytes from progenitor cells. FGF-2 are also involved in a variety of biological processes, including embryonic development, morphogenesis, tissue repair, tumor growth, and invasion. As a multifunctional cytokine, FGF-2 is first isolated from the pituitary. Later, it was identified from various cell types including cardiac myocytes, cardiac fibroblasts, endothelial cells, and smooth muscle cells.

Uniprot ID

NP_001392443

Synonyms

Fibroblast Growth Factors

Sequence Note

Ala2-Ser155

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration of 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 2 weeks under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Image

Tainan Headquarters

+886-6-2536677


bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677



SDS-PAGE analysis of recombinant swine FGF-2.

Disclaimer : For Research Use or Further Manufacturing Only.