

Mouse IL-1 Alpha, His Tag, E. coli

Catalog Number LDG0

Package

LDG022PME

5 μg / 20 μg / 100 μg / Customized package

For full product information, images and publications, please visit our website.



Specifications

Species of Origin	Expression System
Mouse	Escherichia coli
Affinity Tag	Storage Buffer
His Tag (C-term)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Purity	Molecular weight
>95% as determined by SDS-PAGE analysis.	The protein has a calculated MW of 18.93 kDa. The protein migrates as 20 kDa under reducing condition (SDS-PAGE analysis).
Activity	Endotoxin Level
Measure by its ability to induce D10.G4.1 cells proliferation. The ED ₅₀ for this effect is <5 pg/mL. The specific activity of recombinant mouse IL-1 alpha is > 2 x 10 ⁸ IU/mg.	<0.1 EU per 1 μ g of the protein by the LAL method.
Form	

Lyophilized

Background

Tainan Headquarters

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw



Background

Interleukin-1 alpha (IL1 alpha or IL1α) is a member of the interleukin-1 cytokine family, found constitutively present in epithelial layers of the entire gastrointestinal tract, lung, liver, kidney, endothelial cells, and astrocytes. The synthesized IL-1 alpha is a 31 kDa inactive precursor and can be cleaved by intracellular caspase-1 or extracellular proteases to generate the bioactive 17 kDa form and the 16 kDa N-terminal cleavage product. Both precursor and mature IL-1 alpha protein bind to the IL-1 receptor (IL-1R), initiating a cascade of inflammatory cytokines and chemokines production such as IL-6, IL-8, and TNF, in response to viral and bacterial pathogens conditions. IL-1 alpha plays a central role in immune-surveillance mechanisms, stimulating macrophages, neutrophils, and CD8+ T cells activity.

Synonyms

Interleukin-1 alpha, IL-1 alpha

Sequence Note

Ser115-Ser270

Uniprot ID

#P01582

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H_2O to a concentration of 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Tainan Headquarters

Innovation & Research Center

CLD Center

☑ bd@leadgene.com.tw

& +886-2-27065528

& +886-6-2536677



Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Image

kDa	M		
75 60	LEAD	GENE	
45	-		
35	-	Ø	
25	-		
15		GENE	
10			

SDS-PAGE analysis of recombinant mouse IL-1 alpha.

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarters

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-2-27065528

☑ bd@leadgene.com.tw

& +886-6-2536677