

Human FGF-18, His Tag, E. coli

Catalog Number LDG08

Package

LDG082PHE

5 μg / 20 μg / 100 μg / Customized package

For full product information, images and publications, please visit our website.



Specifications

Species of Origin	Expression System
Human	Escherichia coli
Affinity Tag	Storage Buffer
His Tag (C-term)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 8.0.
Purity	Molecular weight
>98% as determined by SDS-PAGE analysis.	The protein has a calculated MW of 21.11 kDa. The protein migrates as 20 kDa under reducing condition (SDS-PAGE analysis).
Activity	Endotoxin Level
Measure by its ability to induce 3T3 cells proliferation. The ED ₅₀ for this effect is 1.3-2.0 ng/mL. The specific activity of recombinant human FGF-18 is > 5 x 10 ⁵ IU/mg.	<0.1 EU per 1 μg of the protein by the LAL method.
Form	

Lyophilized

Background

Tainan Headquarters

Innovation & Research Center

CLD Center

& +886-6-2<u>53</u>6677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw



Background

Fibroblast Growth Factors-18 (FGF-18) is a 24 kDa member of the fibroblast Growth Factors with 207 amino acid residues. FGF-18 is mainly expressed from glandular and luminal cells, cardiomyocytes, breast myoepithelial cells and endometrial ciliated cells. FGF-18 involved cell proliferation, cell differentiation and cell migration. FGF-18 is an important role in skeletal growth and development, stimulates hepatic and intestinal proliferation.

Synonyms

zFGF5, Fibroblast Growth Factors 18

Sequence Note

Ala27-Arg199

Shipping

storage.

The product is shipped with polar packs. Upon receipt,

store it immediately at -20°C or lower for long term

Uniprot ID

#O76093

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration of 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Image

Tainan Headquarters

Innovation & Research Center

CLD Center


& +886-6-2536677

& +886-6-2536677

☑ bd@leadgene.com.tw

& +886-2-27065528





SDS-PAGE analysis of recombinant human FGF-18.

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarters

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw