

## Human RANKL, His Tag, E. coli

<b>Catalog Number</b>	LDG122PHE
<b>Package</b>	5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



### Specifications

#### Species of Origin

Human

#### Affinity Tag

His Tag (C-term)

#### Purity

>98% as determined by SDS-PAGE analysis.

#### Activity

Measure by its ability to induce osteoclast differentiation in RAW264.7 cells. The ED<sub>50</sub> for this effect is <10 ng/mL.

#### Form

Lyophilized

#### Expression System

Escherichia coli

#### Storage Buffer

Lyophilized from a 0.2 µm filtered solution of PBS, pH 8.0.

#### Molecular weight

The protein has a calculated MW of 20.67 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).

#### Endotoxin Level

<0.1 EU per 1 µg of the protein by the LAL method.

### Background

#### Tainan Headquarters

+886-6-2536677

bd@leadgene.com.tw

#### Innovation & Research Center

+886-2-27065528

#### CLD Center

+886-6-2536677

### Background

Receptor activator of NF- $\kappa$ B (RANK) ligand (RANKL) is type II transmembrane protein with an extracellular domain at the carboxy-terminus of TNF cytokine superfamily. RANKL is a 19.8 kDa protein containing 317 residues and high express in T cells and T cell rich organs, such as thymus and lymph nodes. RANKL-RANK (RANKL receptor) plays an important role in bone metabolism, dysregulation, and immune system

### Uniprot ID

#O14788

### Synonyms

Receptor activator of nuclear factor kappa-B ligand, RANKL, Osteoclast differentiation factor, ODF, Osteoprotegerin ligand, OPGL, TNF-related activation-induced cytokine, TRANCE, Tumor necrosis factor ligand superfamily member 11, TNFL11

### Sequence Note

Glu143-Asp317

## Instruction

### Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration of 200  $\mu$ g/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

### Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

### Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

## Image

### Tainan Headquarters

+886-6-2536677


bd@leadgene.com.tw

### Innovation & Research Center

+886-2-27065528

### CLD Center

+886-6-2536677



SDS-PAGE analysis of recombinant human RANKL.

**Disclaimer :** For Research Use or Further Manufacturing Only.