

Human CXCL4, His Tag, E. coli

Catalog Number LDG156PHE

Package 5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Specifications

Species of Origin

Human

Affinity Tag

His Tag (N-term)

Purity

>98% as determined by SDS-PAGE analysis.

Activity

Measure by its ability to inhibit human FGF-2-induce proliferation in HUVEC cells. The ED₅₀ for this effect is <5 µg/mL.

Form

Lyophilized

Expression System

Escherichia coli

Storage Buffer

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Molecular weight

The protein has a calculated MW of 8.58 kDa. The protein migrates as 14 kDa under reducing condition (SDS-PAGE analysis).

Endotoxin Level

<0.1 EU per 1 µg of the protein by the LAL method.

Background

Tainan Headquarter

+886-6-2536677 +886-6-2531536

info@leadgene.com.tw

Taipei Office

+886-2-27065528

Background

C-X-C motif chemokine 4 (CXCL4) also named platelet factor 4 (PF4), which is a chemokine of the intercrine alpha family. CXCL4 is a 8kDa protein containing 70 amino acid residues. CXCL4 is produced by the activated platelets which plays an important role in immune responses. CXCL4 inhibit the cell proliferation, platelet aggregation and wound repair. CXCL4 also suppresses the hematopoiesis.

Uniprot ID

P02776 2

Synonyms

C-X-C motif chemokine 4, Iroplact, Oncostatin-A, Platelet factor 4

Sequence Note

Glu32-Ser101

Instruction**Reconstitution**

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.


Image**Tainan Headquarter**

☎ +886-6-2536677 📠 +886-6-2531536


✉ info@leadgene.com.tw

Taipei Office

☎ +886-2-27065528



Human CXCL4, His Tag, E. coli (LDG156PHE) inhibited human FGF-2-induced HUVEC cell proliferation, with the ED50 at 4.87 µg/mL.



SDS-PAGE analysis of recombinant human CXCL4.

Disclaimer : For Research Use or Further Manufacturing Only.