

Human Heregulin Beta 1, His Tag, E. coli

Catalog Number	LDG185PHE
Package	5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Specifications

Species of Origin

Human

Expression System

Escherichia coli

Affinity Tag

His Tag (C-term)

Storage Buffer

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Purity

>95% as determined by SDS-PAGE analysis.

Molecular weight

The protein has a calculated MW of 8.42 kDa.
The protein migrates as 8 kDa under reducing condition (SDS-PAGE analysis).

Activity

Measure by its ability to induce MCF-7 cells proliferation. The ED₅₀ for this effect is < 10 ng/mL.

Endotoxin Level

<1.0 EU per 1 µg of the protein by the LAL method.

Form

Lyophilized

Background

Tainan Headquarters

+886-6-2536677

bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677

Background

Neuregulin-1 (NRG-1, also called heuregulin1 or neu differentiation factor) is a glycoprotein that belongs to the neuregulins family. Structurally, Neuregulin-1 harbors tissue-specific N terminal sequence, followed by immunoglobulin-like (Ig-like) domains, an EGF-like domain, a transmembrane domain, and a cytoplasmic domain. NRG1 has multiple isoforms produced by alternative splicing. Heregulin- β 1 (HRG- β 1) is one of the isoforms, has been reported to engage the development and survival of cardiomyocytes derived from embryonic stem (ES) cells via activating MAPK-ERK and PI3K-AKT pathways. Moreover, HRG- β 1 plays a central role in promoting the proliferation of neuronal progenitors from embryonic neural stem cells.

Uniprot ID

#Q02297-6

Synonyms

NRG1 Beta 1, Heregulin- β 1

Sequence Note

Ser177-Glu241

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration of 200 μ g/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Tainan Headquarters

+886-6-2536677

bd@leadgene.com.tw


Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677

Image



SDS-PAGE analysis of recombinant human Heregulin beta 1.

Disclaimer : For Research Use or Further Manufacturing Only.