

Product Information & Manual

Information of other products is available at: www.leadgenebio.com

Glycerol Dehydrogenase (GlyDH)

Cat no. LDG0020RG

Product Overview

Description

Glycerol dehydrogenase is an enzyme that catalyzes the oxidation of glycerol to dihydroxyacetone, using NAD⁺ as a cofactor, which is reduced to NADH in the process. This enzyme plays a key role in the metabolism of glycerol and is involved in pathways such as gluconeogenesis and lipid metabolism. Glycerol dehydrogenase is utilized in various industrial applications, including the production of dihydroxyacetone for cosmetic and pharmaceutical products.

Expression system

Escherichia coli

Specification

Appearance	Light brown amorphous powder, lyophilized
Activity	40 U/ mg or more (containing approx. 10% of stabilizers)

Properties

Stability	Stable at -20°C for at least one year
Molecular weight	40.9 kDa
Isoelectric point	5.27

Reconstitution


It is recommended to weight and reconstitute 30 mg of lyophilized powder in 157 µL double-distilled water directly and incubate the solution for at least 10 mins to ensure sufficient re-dissolved.

Applications

1. Enzymatic determination of glycerol
2. Research of energy metabolism⁽¹⁾
3. Glycerol metabolism of pathogenic bacteria⁽¹⁾

Assay

1. Assay principle



2. Unit definition

One unit causes the formation of one micromole of NADH per minute under the conditions described below.

3. Reagents

A. Carbonate-bicarbonate buffer, pH 11.0	0.2 M (Prepare by mixing 0.2 M K ₂ CO ₃ and 0.2 M NaHCO ₃ to reach pH 11.0)
B. Glycerol solution	0.3 M
C. Ammonium sulfate solution	1.0 M
D. NAD ⁺ solution	10 mM [Weigh 133 mg of NAD ⁺ (MW = 663.43) and dissolve in 18 mL of H ₂ O. Adjust the pH to 7.0 with 0.5 N KOH and fill up to 20.0 mL with H ₂ O] (Should be prepared freshly)
E. Enzyme diluent	20 mM K-phosphate buffer pH 7.5

4. Procedure

- Prepare the following working solution immediately before use (for 4 reactions).

Working solution

Carbonate-bicarbonate buffer, pH 11.0 (Reagent A)	3 mL
Glycerol solution (Reagent B)	2.2 mL
Ammonium sulfate solution (Reagent C)	0.2 mL
NAD ⁺ solution (Reagent D)	0.6 mL
Total	6 mL

Be sure the pH in the range (pH 10.0–10.5). If not, adjust the pH to 10.5 with 1.0 N KOH or 1.0 N HCl, and store on ice in a brownish bottle.

- Pipette 1.45 mL of the working solution into a tube and equilibrate at 25°C for about 5 minutes.
- Add 0.05 mL of the enzyme solution* and mix by gentle inversion.

Concentration in a reaction	
Carbonate-bicarbonate buffer	0.1 M
Glycerol	0.1 M
NAD ⁺	1 mM
Ammonium sulfate solution	33 mM

- Add 1 mL of the mixture into a cuvette (d = 1.0 cm).
- Record the increase in optical density at 340 nm against water for 1 to 5 minutes in a spectrophotometer at room temperature and calculate the ΔOD per minute from the initial linear portion of the curve (ΔOD test). At the same time, measure the blank rate (ΔOD blank) by using the same method as the test except that the enzyme diluent is added instead of enzyme solution.

* Dilute the enzyme in ice-cold enzyme diluent (Reagent E) to 0.10–0.25 U/ mL and store on ice.

- Activity can be calculated by using the following formula:

Volume activity (U/mL) =

$$\frac{\Delta OD / \text{min} (\Delta OD \text{ test} - \Delta OD \text{ blank}) \times V_t \times df}{6.22 \times 1.0 \times V_s}$$

$$= \Delta OD / \text{min} \times 4.82 \times df$$

Weight activity (U/mg) = (U/mL) × 1/C

V_t: Total volume (1.5 mL)

V_s: Sample volume (0.05 mL)

6.22 : Millimolar extinction coefficient of NADH (cm²/micromole)

1.0: Light path length (cm)

df: Dilution factor

C: Enzyme concentration in dissolution (mg/mL)

Reference

- Cedric Blötz. et al.* Glycerol metabolism and its implication in virulence in *Mycoplasma*. *FEMS Microbiology Reviews* (2017).

The effect of different conditions on Glycerol dehydrogenase

A.




Figure A. Temperature activity of Glycerol dehydrogenase. The enzyme reactions in 0.1 M Carbonate-bicarbonate buffer, pH 11.0, were carried out under different temperature.

B.




Figure B. pH activity of Glycerol dehydrogenase. The buffer conditions with various pH values were used in the reaction at 25°C. pH 4.0-6.0, 0.1 M Sodium citrate buffer; pH 7.5-8.0, 0.1 M Potassium phosphate buffer; pH 10.0-11.0, 0.1 M Carbonate-bicarbonate buffer.

C.




Figure C. Thermal stability of Glycerol dehydrogenase. The enzyme powder was reconstituted by double-distilled water and treated with different temperatures for 15 minutes. Final concentration: 16 U/ mL

D.




Figure D. pH stability of Glycerol dehydrogenase. The enzyme powder was reconstituted by double-distilled water and treated with different pH buffer condition for 20 hours at 25°C. pH 6.0, 0.1 M Sodium citrate buffer; pH 7.5-8.0, 0.1 M Potassium phosphate buffer; pH 10.0-11.0, 0.1 M Carbonate-bicarbonate buffer.

Disclaimer

For Research Use or Further Manufacturing Only.