

Streptavidin-HRP

Catalog Number	LDG0002RO	infor publ
Package	500 μL / Customized package	our
O Publications (1		

For full product information, images and publications, please visit our website.



Overview

)

Description

Streptavidin is a 53 kDa homotetramer isolated from Streptomyces avidinii. Because of its strong non-covalent interaction with biotin, streptavidin can be used to detect and isolate biotinylated proteins. Conjugation of horseradish peroxidase (HRP) to streptavidin is obtained by cross linking the amino groups on streptavidin with the carbohydrate groups on HRP. Streptavidin-HRP is typically used biotinylated antibodies or proteins.

Product Note

Recommended dilution factor: ELISA: 1:5000-20000 WB: 1:1000-10000 IHC: 1:200-500 Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions. Horseradish peroxidase is conjugated to the streptavidin tetramer at about 1:1 molar ratio. Optimal titers for the applications should be determined by the researcher.

Specifications	
Application ELISA, WB, IHC	Conjugation
Buffer	Form
Liquid. Phosphate Buffered Saline containing 50% glycerol and 0.01% ProClin 300, pH 7.4.	Liquid

Tainan Headquarter	Innovation & Research Center	CLD Center
𝔅 +886-6-2536677		& +886-6-2536677
☑ bd@leadgene.com.tw		



Instruction

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage

This product is stable after storage at:

- 2-8°C for 6 months under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles. Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw