

回走際周

For full product

# **HRV 3C Protease**

Catalog Number	LDG0009RG	information, images and publications, please visit
Package	Customized package / 1,000 U / 10,000 U	our website.
66 Publications (1		

)

# Overview

## Description

HRV 3C Protease is a recombinant form of the 3C protease derived from human rhinovirus 14 expressed in E. coli (specific activity 1800-2000 U/mg). This product is a highly purified recombinant 6XHis-fusion protein. This protease requires neither metal nor cofactors for activity. HRV 3C Protease recognizes the cleavage site: Leu-Glu-Val-Leu-Phe-Gln1Gly-Pro (LEVLFQ1GP).

## **Product Note**

- Cleavage efficiency may differ based on structure and properties of each target protein, we recommend testing several enzyme-to-substrate ratios, temperatures and incubation times.
- HRV 3C Protease reactions can be performed in a buffer which is optimal for the target protein. Reducing reagents (e.g., DTT) or salts (e.g., NaCl) can be added for cleavage efficiency evaluation.

#### Components

Package	Items	Quantity
1.000 U	HRV 3C Protease	1 vial (1,000 U)
1,000 0	10X HRV 3C Cleavage Buffer	1 vial (10 mL)
10,000 U	HRV 3C Protease	1 vial (10,000 U)
10,000 0	10X HRV 3C Cleavage Buffer	1 vial (10 mL)

# **Specifications**

Tainan Headquarter	Innovation & Research Center	CLD Center
& +886-6-2536677	& +886-2-27065528	& +886-6-2536677
☑ bd@leadgene.com.tw		



## **Expression system**

## Escherichia coli

#### Purity

>98% as determined by SDS-PAGE analysis.

<1 EU per 1 µg of the protein by the LAL method.

#### **Buffer**

Lyophilized from a solution containing 50 mM Tris, 150 mM NaCl, 1 mM EDTA, 1 mM DTT, 0.04% Tween20, 8% trehalose, 8% mannitol.

#### **Unit Definition**

One unit of HRV 3C Protease is defined as the amount of enzyme that will cleave >95% of 0.1 mg HRV 3C cleavage control protein in 150 mM NaCl, 50 mM Tris-HCl pH 7.5, at 4°C for 16 h.

The product is shipped with polar packs. Upon receipt,

store it immediately at -20°C or lower for long term

#### Form

Lyophilized

Shipping

storage.

# Instruction

Endotoxin level

#### Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

#### Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

# Image

#### **Tainan Headquarter**

**Innovation & Research Center** 

# **CLD** Center


& +886-6-2536677


& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw







HRV 3C protease removes the purification tag appended to interest proteins after the affinity chromatography step. SDS-PAGE analysis of recombinant HRV 3C Protease Protein

Disclaimer : For Research Use or Further Manufacturing Only.

## **Tainan Headquarter**

Innovation & Research Center

**CLD** Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw