

LeadGMP® Human IL-2 Protein, His Tag, E. coli

Catalog Number LDG003PHE-GMP

Package

100 μg / 1 mg / Customized package

For full product information, images and publications, please visit our website.



Specifications

| Species of Origin | Expression system |
|---|--|
| Human | Escherichia coli |
| Affinity Tag | Buffer |
| | |
| His Tag (C-term) | Lyophilized from a 0.2 μ m filtered solution of NaPi buffer, 0.018% SDS, pH 7.5. |
| Purity | Molecular weight |
| >97% as determined by SDS-PAGE analysis. | The protein has a calculated MW of 16.4 kDa. |
| | The protein migrates as 13 kDa under reducing |
| | condition (SDS-PAGE analysis). |
| | |
| Activity | Endotoxin level |
| Measure by its ability to induce proliferation in CTLL-2 | <0.05 EU per 1 μ g of the protein by the LAL method. |
| cells. The ED $_{50}$ for this effect is <3 ng/mL. The specific | |
| activity of recombinant human IL-2 is >5 x 10 ⁶ IU/mg, | |
| which is calibrated against the human IL-2 WHO | |
| International Standard (NIBSC code: 86/500). | |
| | |
| Mycoplasma | Form |
| Not detected | Lyophilized |
| | |
| | |

Background

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677



Background

Interleukin-2 (IL-2) is a 15.5 kDa four bundle, immunomodulatory cytokine which is expressed predominately by antigen-simulated CD4+ T cells, CD8+ T cells, natural killer (NK) cells, and dendritic cells (DC). IL-2 plays a fundamental role in promoting NK cell proliferation, cytotoxicity, and B cells differentiation. IL-2 modulate T cell differentiation programs, promoting naïve CD4+ T cell differentiation into T helper-1 (Th1) and T helper-2 (Th2) cells.

Synonyms

Interleukin-2, T-cell growth factor , TCGF

Sequence Note

Ala21-Thr153

Uniprot ID

P60568

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H_2O to a concentration not less than 0.5 mg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-6-2536677



Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.


Avoid repeated freeze/thaw cycles.

Manufacturing specifications


LeadGMP® recombinant proteins are manufactured in ISO 13485:2016 and GMP certified facility. The processes include:

- Animal-free reagent and laboratory
- Manufactured and tested under GMP guideline
- Testing and traceability of raw material
- Records of the maintenance and equipment calibration
- Personnel training records
- Batch-to-batch consistency
- Documentation of QA control and process changes
- Manufactured and tested under an ISO 13485:2016 certified quality management system
- Stability monitor of product shelf-life


Image



The purity of LeadGMP® Human IL-2 Protein is greater than 95% as determined by SEC-HPLC.



LeadGMP® Human IL-2 Protein, His Tag, E. coli (LDG003PHE-GMP) induces CTLL-2 cell proliferation, with the ED50 < 3.0 ng/mL.



SDS-PAGE analysis of LeadGMP® IL-2, Human

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2536677

® +886-2-27065528

& +886-6-2536677



Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677