

Anti-c-Met Antibody [Clone 4A9]

Catalog Number	LDG0045YA
Package	100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Overview

Description

The c-mesenchymal epithelial transition factor (c-Met ; also known as hepatocyte growth factor receptor, HGFR), is a receptor tyrosine kinase (RTK) that mainly exists in epithelial cells. c-Met and its high-affinity ligand, hepatocyte growth factor (HGF), play important roles in mediating embryogenesis, tissue regeneration, wound healing and the formation of nerve and muscle. Aberrant HGF/c-Met axis activation is associated with the proliferation, survival, invasion and metastasis of various tumor cells, and thus c-Met may be a tumor biomarker and therapeutic target.

Product Note

Recommended dilution factor:

ELISA: 1:5000-20000

WB: 1:1000-10000

IFA: 1:200-1000

FACS: Assay dependent

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

Specifications

Host	Clonality
Mouse	Monoclonal
Isotype	Clone Name
IgG1	clone 4A9
Immunogen	Reactivity
c-Met	Human

Tainan Headquarter

+886-6-2536677

bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677

Application

ELISA, WB, IFA, FACS

Concentration

1 mg/mL

Specificity

c-Met

Conjugation

Unconjugated

Buffer

Phosphate Buffered Saline containing 0.03% ProClin 300, pH 7.4.

Form

Liquid

Instruction**Shipping**

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.



Stability & Storage


This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.


Avoid repeated freeze/thaw cycles.

Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.


Image**Tainan Headquarter** +886-6-2536677 bd@leadgene.com.tw**Innovation & Research Center** +886-2-27065528**CLD Center** +886-6-2536677




Immunofluorescence analysis of Anti-c-Met Antibody [Clone 4A9]
 MCF7 cells were fixed in 100% methanol, permeabilized with PBS containing 0.1% Triton X-100. Cells were stained with mouse anti-c-Met monoclonal antibody (1:200) followed by secondary antibodies (goat anti-Mouse IgG-iFluor 488, 1:200, green) and cell nuclei were stained with Hoechst 33342 (Blue).



FACS analysis of Anti-c-Met Antibody [Clone 4A9]
 MCF-7 cells were stained with anti-c-Met monoclonal antibody at 2 µg/ml (red) and without antibody control (black).



ELISA titration of Anti-c-Met Antibody [Clone 4A9]
 Titration curve of anti-c-Met antibody in ELISA. Red: c-Met; Black: BSA (negative control).



Western blotting analysis of Anti-c-Met Antibody [Clone 4A9]
 HeLa cell lysates (50 µg) were stained with mouse anti-c-Met monoclonal antibody at 1:5000 dilution.

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarter

Innovation & Research Center

CLD Center

+886-6-2536677

+886-2-27065528

+886-6-2536677

bd@leadgene.com.tw

Tainan Headquarter

☎ +886-6-2536677

✉ bd@leadgene.com.tw

Innovation & Research Center

☎ +886-2-27065528

CLD Center

☎ +886-6-2536677