

Donkey Anti-Goat IgG (H+L) Antibody (TAMRA)

LDG0048YE **Catalog Number**

Package

500 µL / 1 mL / Customized package

For full product information, images and publications, please visit our website.



Overview

Description

TAMARA is a fluorescent dye for DNA, peptides and proteins. Leadgene Donkey Anti-Goat IgG (H+L) Antibody (TAMRA) is conjugated with high-activity TAMARA

Product Note

Recommended dilution factor: IFA: 1:200-1000 FACS: 1:200-1000

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

Host	Clonality
Donkey	Polyclonal
Immunogen	Reactivity
Purified goat IgG	Goat (Minor cross-reactivity to human in higher concentration)
Application	Conjugation
IFA, FACS	5(6)-TAMRA
Concentration	Buffer
0.5 mg/mL	Phosphate Buffered Saline containing 1% BSA, 0.03% ProClin 300, pH 7.4.

Tainan Headquarter

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw



Specificity IgG	Form Liquid
Instruction	
Shipping The product is shipped with polar packs. Upon receipt, store it immediately at 2-8°C for long term storage.	 Stability & Storage This product is stable after storage at: 2-8°C for 6 months under sterile conditions from date of receipt. Do not freeze and protect from light.

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw