

## Human IL-2, His Tag, E. coli

<b>Catalog Number</b>	LDG004PHE
<b>Package</b>	5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



### Specifications

#### Species of Origin

Human

#### Affinity Tag

His Tag (C-term)

#### Purity

>98% as determined by SDS-PAGE analysis.

#### Activity

Measure by its ability to induce proliferation in CTLL-2 cells. The ED<sub>50</sub> for this effect is <3 ng/mL. The specific activity of recombinant human IL-2 is >5 x 10<sup>6</sup> IU/mg.

#### Form

Lyophilized

#### Expression system

Escherichia coli

#### Buffer

Lyophilized from a 0.2 µm filtered solution of NaPi buffer, 0.018% SDS, pH 7.5.

#### Molecular weight

The protein has a calculated MW of 16.4 kDa. The protein migrates as 13 kDa under reducing condition (SDS-PAGE analysis).

#### Endotoxin level

<0.05 EU per 1 µg of the protein by the LAL method.

### Background

#### Tainan Headquarter

+886-6-2536677

bd@leadgene.com.tw

#### Innovation & Research Center

+886-2-27065528

#### CLD Center

+886-6-2536677

### Background

Interleukin-2 (IL-2) is a 15.5 kDa four bundle, immuno-modulatory cytokine which is expressed predominately by antigen-simulated CD4+ T cells, CD8+ T cells, natural killer (NK) cells, and dendritic cells (DC). IL-2 plays a fundamental role in promoting NK cell proliferation, cytotoxicity, and B cells differentiation. IL-2 modulate T cell differentiation programs, promoting naïve CD4+ T cell differentiation into T helper-1 (Th1) and T helper-2 (Th2) cells.

### Uniprot ID

#P60568

### Synonyms

Interleukin-2, T-cell Growth Factors , TCGF

### Sequence Note

Ala21-Thr153

## Instruction

### Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration of 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

### Stability & Storage

This product is stable after storage at:


- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.


### Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

## Image



The purity of recombinant human IL-2 is greater than 95% as determined by SEC-HPLC.



SDS-PAGE analysis of recombinant human IL-2

**Disclaimer :** For Research Use or Further Manufacturing Only.