

# LeadGMP® PNGase F

Catalog Number

LDG004R-GMP

Package

15 KU / 75 KU / Customized package

For full product information, images and publications, please visit our website.



## **Overview**

#### Description

PNGase F is an enzyme used in biochemistry and molecular biology to remove N-linked glycans from glycoproteins. By using PNGase F, researchers can enzymatically cleave between these sugar chains and asparagine residues of glycoproteins, allowing for the study of protein structure and function, particularly in glycosylation research.

#### Components

Package	Items	Quantity
	LeadGMP® PNGase F1 vial (400 U/µL)10X Glycoprotein Denaturing Buffer1 vial (1 mL)10X Reaction buffer1 vial (1 mL)10% NP-401 vial (1 mL)	1 vial (400 U/µL)
15 KU		1 vial (1 mL)
		1 vial (1 mL)
		1 vial (1 mL)
	LeadGMP® PNGase F	PNGase F 1 vial (400 U/µL)
75 KU	10X Glycoprotein Denaturing Buffer	1 vial (1 mL)
	10X Reaction buffer 1 vial (1 mL)	1 vial (1 mL)
	10% NP-40	1 vial (1 mL)

# **Specifications**

Tainan Headquarter	Innovation & Research Center	CLD Center
& +886-6-2536677	& +886-2-27065528	<u></u>
☑ bd@leadgene.com.tw		



#### Expression system

## Escherichia coli

#### Buffer

20mM Tris-HCI, 50mM NaCl, 5mM EDTA, pH 7.5

## **Unit Definition**

One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 10  $\mu$ g of denatured RNase B in 1 hour at 37°C in a total reaction volume of 20  $\mu$ L.

#### Mycoplasma

Not detected

Concentration

400 U /µL

## Purity

>95% as determined by SDS-PAGE analysis.

## **Endotoxin level**

<0.05 EU per 1  $\mu$ g of the protein by the LAL method.

Form

Liquid

# Instruction

## Tainan Headquarter

Innovation & Research Center

**CLD** Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw



#### Stability & Storage

This product is stable after storage at:


 -20°C for long-term storage under sterile conditions. Avoid repeated free-thaw cycles.

#### Manufacturing specifications


LeadGMP® recombinant proteins are manufactured in ISO 13485:2016 and GMP certified facility. The processes include:

- Animal-free reagent and laboratory
- Manufactured and tested under GMP guideline
- Testing and traceability of raw material
- Records of the maintenance and equipment calibration
- Personnel training records
- Batch-to-batch consistency
- Documentation of QA control and process changes
- Manufactured and tested under an ISO 13485:2016 certified quality management system
- Stability monitor of product shelf-life


# Image



The intracellular PD-L1 protein was digested with LeadGMP® PNGase F and analyzed by Western blotting. The assay was performed by incubating 400 U PNGase F and 20 µg of glycoprotein from whole cell lysate.



SDS-PAGE analysis of recombinant PNGase F



PNGase F cleaves high mannose, hybrid-type and complex structure glycans.

## Tainan Headquarter

Innovation & Research Center

## **CLD** Center

**⊗** +886-6-2536677

*ⓑ* +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw



Disclaimer : For Research Use or Further Manufacturing Only.

# Tainan Headquarter

Innovation & Research Center

**CLD** Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw