

Anti-SARS-CoV & CoV-2 NP Antibody [Clone 103]

Catalog Number LDG01

Package

LDG0135YA

100 µg / 1 mg / Customized package

For full product information, images and publications, please visit our website.



Overview

Description

Human anti-SARS-CoV & CoV-2 NP antibody recognize human SARS Coronavirus Nucleocapsid protein (NP). Coronavirus NP localize to the cytoplasm and the nucleolus in both Virus-like particle (VLP) infected primary cells and in cells transfected with NP plasmid. NP has abundant expression in coronavirus and is a highly immunogenic phosphoprotein. NP is also conserved in sequence. Due to these characteristics above, the NP is an ideal marker for diagnosis.

Product Note

Recommended dilution factor: ELISA: 1:5000-20000

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

Specifications

Clonality	Isotype
Recombinant Human IgG	IgG1
Clone Name	Reactivity
clone 103	SARS-CoV & CoV-2
Application	Conjugation
ELISA, CLIA, LFIA	Unconjugated
Concentration 1 mg/mL	Buffer Phosphate Buffered Saline containing 0.2% ProClin 300, pH 7.4.

Tainan Headquarter Innovation & Research Center CLD Center © +886-6-2536677 © +886-2-27065528 © +886-6-2536677

☑ bd@leadgene.com.tw



Specificity

Nucleocapsid protein

Instruction

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage

Form

Liquid

This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles. Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.

Image

3-	- Omicron NP
	SARS-CoV-2 N
1- 0 200 Human anti-SAR	800 1000 NP Antibody

ELISA titration of Human anti-SARS-CoV & CoV-2 NP Antibody, clone 103

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarter

Innovation & Research Center

CLD Center

<u>®</u> +886-6-253<u>6</u>677

® +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw