

# Anti-CD28 Chimeric IgG Antibody [Clone 0AA02]

**Catalog Number** LDG0179YA **Package** 100 µg / Customized package

For full product information, images and publications, please visit our website.



## Overview

## Description

CD28, a well-known costimulatory receptor expressed on T cells, is a type I transmembrane glycoprotein of the Ig superfamily. When the T cells receive the first signal delivered by APCs (antigen-presenting cells), it initiates signaling cascades, which favor T cell survival and proliferation. It has been demonstrated that patients with lung cancer who responded to PD-1 therapy had more CD28+ T cells, suggesting that CD28 may predict treatment response.

**Product Note** 

Recommended dilution factor:

ELISA: 1:5000-20000

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

# **Specifications**

Clonality Recombinant Human IgG **Clone Name** clone 0AA02 **Application ELISA** Concentration 0.5 mg/mL

Isotype

lgG1

Reactivity

Human

Conjugation

Unconjugated

**Buffer** 

Phosphate Buffered Saline containing 0.03% ProClin 300, pH 7.4.

Tainan Headquarter

**Innovation & Research Center** 

**CLD Center** 



**Specificity** 

CD28

**Form** 

Liquid

## Instruction

## **Shipping**

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

## Stability & Storage


This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles.

Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.

## **Image**



ELISA titration of Anti-CD28 Antibody [Clone 0AA02] Titration curve of anti-CD28 antibody in ELISA.

**Disclaimer:** For Research Use or Further Manufacturing Only.

Tainan Headquarter

**Innovation & Research Center** 

**CLD Center**