

Human IL-20, His Tag, E. coli

 Catalog Number
 LDG024PHE

 Package
 5 μg / 20 μg / 100 μg / Customized package

For full product information, images and publications, please visit our website.



Specifications

Species of Origin

Human

Affinity Tag

His Tag (C-term)

Purity

>98% as determined by SDS-PAGE analysis.

Activity

Measure by its ability to chemoattract BaF3 cells transfected with human IL-20R alpha and IL-20R beta. The ED $_{50}$ for this effect is <0.2 ng/mL.

Form

Lyophilized

Expression system

Escherichia coli

Buffer

Lyophilized from a 0.2 μm filtered solution of PBS, pH 8.0

Molecular weight

The protein has a calculated MW of 18.47 kDa. The protein migrates as 16 kDa under reducing condition (SDS-PAGE analysis).

Endotoxin level

<0.1 EU per 1 μg of the protein by the LAL method.

Background



Background

Interleukin 20 (IL-20) predicts a molecular mass of 17.6 kDa. The IL-20 subfamily of cytokines is comprised of IL-19, IL-20, IL-22, IL-24, and IL-26. It regulates of differentiation of keratinocytes during inflammation, the expansion of multipotential hematopoietic progenitor cells.

Uniprot ID

#Q9NYY1

Synonyms

Cytokine Zcyto10, Interleukin-20

Sequence Note

Leu25-Glu176

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H_2O to a concentration of 200 $\mu g/mL$ and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.


Avoid repeated freeze/thaw cycles.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Image





SDS-PAGE analysis of recombinant human IL-20.

Disclaimer : For Research Use or Further Manufacturing Only.