

Mouse IL-10, His Tag, E. coli

Catalog Number LDG03

Package

LDG032PME

5 μg / 20 μg / 100 μg / Customized package

For full product information, images and publications, please visit our website.



Specifications

Species of Origin	Expression system
Mouse	Escherichia coli
Affinity Tag	Buffer
His Tag (C-term)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Purity	Molecular weight
>98% as determined by SDS-PAGE analysis.	The protein has a calculated MW of 19.7 kDa. The protein migrates as 19 kDa under reducing condition (SDS-PAGE analysis).
Activity	Endotoxin level
Measure by its ability to induce MC/9-2 cells proliferation. The ED ₅₀ for this effect is <1 ng/mL. The specific activity of recombinant mouse IL-10 is > 1 x 10^6 IU/mg.	<0.1 EU per 1 μg of the protein by the LAL method.
Form	

Lyophilized

Background

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2<u>53</u>6677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw



Background

Interleukin-10 (IL-10), also known as cytokine synthesis inhibitory factor (CSIF), is an anti-inflammatory cytokine. Many different types of cells can produce IL-10, including immune cells and non-immune cells. IL-10 exerts inhibitory functions on DCs and macrophages, which is a potent inhibitor of antigen presentation and limit the production of the Th1-associated cytokines IL-2 and interferon-y (IFN-y). IL-10 is also a key immunoregulator during infection due to its inhibitory effect on inflammatory cytokine production. Consequently, the excessive Th1 and CD8+ T cell responses could be suppressed during infection.

Uniprot ID

#P18893

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration of 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Synonyms

Interleukin-10, Cytokine synthesis inhibitory factor, CSIF

Sequence Note

Ser19-Ser178

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Image

Tainan Headquarter

Innovation & Research Center

CLD Center


& +886-6-2536677

& +886-2-27065528

☑ bd@leadgene.com.tw

& +886-6-2536677





SDS-PAGE analysis of recombinant mouse IL-10.

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw