

## Human FGF-1, His Tag, E. coli

**Catalog Number** LDG067PHE

**Package** 5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



### Specifications

#### Species of Origin

Human

#### Affinity Tag

His Tag (C-term)

#### Purity

>98% as determined by SDS-PAGE analysis.

#### Activity

Measure by its ability to induce 3T3 cells proliferation. The ED<sub>50</sub> for this effect is <0.3 ng/mL. The specific activity of recombinant human FGF-1 is > 1 x 10<sup>6</sup> IU/mg.

#### Form

Lyophilized

#### Expression system

Escherichia coli

#### Buffer

Lyophilized from a 0.2 µm filtered solution of PBS, pH 8.0.

#### Molecular weight

The protein has a calculated MW of 16.77 kDa. The protein migrates as 18kDa under reducing condition (SDS-PAGE analysis).

#### Endotoxin level

<0.1 EU per 1 µg of the protein by the LAL method.

### Background

#### Tainan Headquarter

+886-6-2536677 +886-6-2531536

info@leadgene.com.tw

#### Taipei Office

+886-2-27065528

### Background

Fibroblast Growth Factors-1 (FGF-1) is a Growth Factors of mitogenic peptides which is a 15 kDa protein containing 139 amino acid residues. Fibroblast Growth Factors-1 is secreted by the macrophage which induce endothelial cell proliferation and angiogenesis. During wound injury, fibroblast Growth Factors-1 helps the tissue remodeling. When the fibroblast Growth Factors-1 binds with the FGFR on the cell membrane which is endocytosed by FGFR and induces the cell cycle.

### Uniprot ID

#P05230

### Synonyms

Acidic fibroblast Growth Factors , aFGF, Endothelial cell Growth Factors , ECGF, Heparin-binding Growth Factors 1 , HBGF-1, Fibroblast Growth Factors 1

### Sequence Note

Phe16-Asp155

## Instruction

### Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

### Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

### Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

## Image


### Tainan Headquarter

+886-6-2536677 +886-6-2531536

info@leadgene.com.tw

### Taipei Office

+886-2-27065528



SDS-PAGE analysis of  
recombinant human FGF-1.

**Disclaimer :** For Research Use or Further Manufacturing Only.

**Tainan Headquarter**

+886-6-2536677 +886-6-2531536

info@leadgene.com.tw

**Taipei Office**

+886-2-27065528