

Human FGF-3, His Tag, E. coli

Catalog Number LDG069PHE

Package 5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Specifications

Species of Origin

Human

Affinity Tag

His Tag (C-term)

Purity

>95% as determined by SDS-PAGE analysis.

Activity

Measure by its ability to induce 3T3 cells proliferation. The ED₅₀ for this effect is <78 ng/mL.

Form

Lyophilized

Expression system

Escherichia coli

Buffer

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Molecular weight

The protein has a calculated MW of 21.99 kDa. The protein migrates as 22 kDa under reducing condition (SDS-PAGE analysis).

Endotoxin level

<0.1 EU per 1 µg of the protein by the LAL method.

Background

Tainan Headquarter

+886-6-2536677 +886-6-2531536

info@leadgene.com.tw

Taipei Office

+886-2-27065528

Background

Fibroblast Growth Factors-3 (FGF-3) also known as INT-2 proto-oncogene, is a 26.9 kDa member of the fibroblast Growth Factors with 239 amino acid residues. FGF-3 regulates embryonic development, cell proliferation and cell differentiation.

Uniprot ID

#P11487

Synonyms

Fibroblast Growth Factors receptor 3, FGFR-3, CD333

Sequence Note

Asp28-Arg212

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 2 weeks under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Image


Tainan Headquarter

☎ +886-6-2536677 📠 +886-6-2531536


✉ info@leadgene.com.tw

Taipei Office

☎ +886-2-27065528



Human FGF-3, His Tag, E. coli (LDG069PHE) induced 3T3 cell proliferation, with the ED₅₀ at 77.51 ng/mL.



SDS-PAGE analysis of recombinant human FGF-3.

Disclaimer : For Research Use or Further Manufacturing Only.