

Mouse Flt-3 Ligand, His Tag, E. coli

Catalog Number	LDG073PME
Package	5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Specifications

Species of Origin

Mouse

Affinity Tag

His Tag (C-term)

Purity

>98% as determined by SDS-PAGE analysis.

Activity

Measure by its ability to induce proliferation in BaF3 cells transfected with mouse Flt-3. The ED₅₀ for this effect is <2 ng/mL.

Form

Lyophilized

Expression system

Escherichia coli

Buffer

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Molecular weight

The protein has a calculated MW of 19.3 kDa. The protein migrates as 19 kDa under reducing condition (SDS-PAGE analysis).

Endotoxin level

<0.1 EU per 1 µg of the protein by the LAL method.

Background

Tainan Headquarter

+886-6-2536677

bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677

Background

Fms-related tyrosine kinase-3 ligand (Flt-3 Ligand) is a protein which is encoded by the Flt3lg gene in mouse. As an important regulator of hematopoiesis, Flt-3 ligand has a tyrosine-protein kinase activity that stimulate the proliferation of hematopoietic progenitor cells of both lymphoid and myeloid origin. Flt-3 ligand can synergistically increase the proliferation of immature progenitors with other cytokines such as GM-CSF, G-CSF, IL-3, EPO, IL-11, IL-12, IL-6 or TPO. Flt-3 ligand is expressed by a variety of hematopoietic progenitor cells including immature hematopoietic and marrow stromal cells.

Uniprot ID

#P49772

Synonyms

Fms-related tyrosine kinase 3 ligand, Flt3L, SL cytokine

Sequence Note

Gly27-Arg188

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration of 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Image

Tainan Headquarter

+886-6-2536677


bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677



SDS-PAGE analysis of recombinant mouse Flt-3 Ligand.

Disclaimer : For Research Use or Further Manufacturing Only.