

Human FGF-8b, His Tag, E. coli

LDG074PHE **Catalog Number Package** $5~\mu g$ / $20~\mu g$ / $100~\mu g$ / Customized package

For full product information, images and publications, please visit our website.



Specifications

Species of Origin

Human

Affinity Tag

His Tag (C-term)

Purity

>95% as determined by SDS-PAGE analysis.

Activity

Measure by its ability to induce 3T3 cells proliferation. The ED50 for this effect is 1.4-3.8 ng/mL. The specific activity of recombinant human FGF-8b is > 2 x 105 IU/mg.

Form

Lyophilized

Expression system

Escherichia coli

Buffer

Lyophilized from a 0.2 µm filtered solution of PBS, pH

Molecular weight

The protein has a calculated MW of 22.14 kDa. The protein migrates as 23 kDa under reducing condition (SDS-PAGE analysis).

Endotoxin level

<0.1 EU per 1 μ g of the protein by the LAL method.

Background

Tainan Headquarter

Innovation & Research Center

CLD Center



Background

Fibroblast Growth Factors-8b (FGF-8b) is a 21.3 kDa member of the fibroblast Growth Factors with 183 amino acid residues. FGF-8 is an important role in the regulation of embryonic development, cell proliferation, cell differentiation and cell migration. In embryogenesis function, FGF8 is necessary in limb, brain, eye and ear formation. FGF-8b having a more potent inductive activity due to higher affinity for Fgf receptors.

Uniprot ID

#P55075-3

Synonyms

Androgen-induced Growth Factors, AIGF, Heparinbinding Growth Factors 8, HBGF-8, Fibroblast Growth Factors 8

Sequence Note

Gln23 - Arg215

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Shipping


The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Image


Tainan Headquarter

Innovation & Research Center

CLD Center



Human FGF-8b, His Tag, E. coli (LDG074PHE) induced 3T3 cell proliferation, with the ED50 at 3.51 ng/mL.



SDS-PAGE analysis of recombinant human FGF-8b.

Disclaimer: For Research Use or Further Manufacturing Only.