

## Human IL-5, His Tag, CHO

<b>Catalog Number</b>	LDG074PHM
<b>Package</b>	5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



### Specifications

#### Species of Origin

Human

#### Expression system

CHO

#### Affinity Tag

His Tag (N-term)

#### Buffer

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

#### Purity

>95% as determined by SDS-PAGE analysis.

#### Molecular weight

The protein has a calculated MW of 13.97 kDa. The protein migrates as 12-22 kDa under reducing condition (SDS-PAGE analysis).

#### Activity

Measure by its ability to induce TF-1 cells proliferation. The ED<sub>50</sub> for this effect is <1 ng/mL.

#### Endotoxin level

<0.1 EU per 1 µg of the protein by the LAL method.

#### Mycoplasma

Not detected.

#### Form

Lyophilized

### Background

#### Tainan Headquarter

+886-6-2536677

bd@leadgene.com.tw

#### Innovation & Research Center

+886-2-27065528

#### CLD Center

+886-6-2536677

### Background

Interleukin-5 (IL-5) is a cytokine that plays a central role in the growth, differentiation, and activation of eosinophils. It is crucial in the pathogenesis of allergic diseases such as asthma and is a target for therapeutic interventions.

### Uniprot ID

P05113

### Synonyms

Interleukin-5, B-cell differentiation factor I, Eosinophil differentiation factor, T-cell replacing factor, TRF

### Sequence Note

Met1-Ser134

## Instruction

### Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

### Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

### Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

## Image

### Tainan Headquarter

+886-6-2536677


bd@leadgene.com.tw

### Innovation & Research Center

+886-2-27065528


### CLD Center

+886-6-2536677



Measure by its ability to induce TF-1 cells proliferation. The ED<sub>50</sub> for this effect is <1 ng/mL.

Activity analysis of recombinant Human IL-5, His Tag, CHO.



SDS-PAGE analysis of recombinant human IL-5.

**Disclaimer :** For Research Use or Further Manufacturing Only.