

## Human FGF-18, His Tag, E. coli

<b>Catalog Number</b>	LDG082PHE
<b>Package</b>	5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



### Specifications

#### Species of Origin

Human

#### Affinity Tag

His Tag (C-term)

#### Purity

>98% as determined by SDS-PAGE analysis.

#### Activity

Measure by its ability to induce 3T3 cells proliferation. The ED<sub>50</sub> for this effect is 1.3-2.0 ng/mL. The specific activity of recombinant human FGF-18 is > 5 x 10<sup>5</sup> IU/mg.

#### Form

Lyophilized

#### Expression system

Escherichia coli

#### Buffer

Lyophilized from a 0.2 µm filtered solution of PBS, pH 8.0.

#### Molecular weight

The protein has a calculated MW of 21.11 kDa. The protein migrates as 20 kDa under reducing condition (SDS-PAGE analysis).

#### Endotoxin level

<0.1 EU per 1 µg of the protein by the LAL method.

### Background

#### Tainan Headquarter

+886-6-2536677

bd@leadgene.com.tw

#### Innovation & Research Center

+886-2-27065528

#### CLD Center

+886-6-2536677

### Background

Fibroblast Growth Factors-18 (FGF-18) is a 24 kDa member of the fibroblast Growth Factors with 207 amino acid residues. FGF-18 is mainly expressed from glandular and luminal cells, cardiomyocytes, breast myoepithelial cells and endometrial ciliated cells. FGF-18 involved cell proliferation, cell differentiation and cell migration. FGF-18 is an important role in skeletal growth and development, stimulates hepatic and intestinal proliferation.

### Uniprot ID

#O76093

### Synonyms

zFGF5, Fibroblast Growth Factors 18

### Sequence Note

Ala27-Arg199

## Instruction

### Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

### Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

### Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

## Image

### Tainan Headquarter

+886-6-2536677


bd@leadgene.com.tw

### Innovation & Research Center

+886-2-27065528

### CLD Center

+886-6-2536677



SDS-PAGE analysis of recombinant human FGF-18.

**Disclaimer :** For Research Use or Further Manufacturing Only.