

Human BMP-16, His Tag, E. coli

Catalog Number LDG1

Package

LDG114PHE

5 μg / 20 μg / 100 μg / Customized package

For full product information, images and publications, please visit our website.



Specifications

Species of Origin	Expression system
Human	Escherichia coli
Affinity Tag	Buffer
His Tag (C-term)	Lyophilized from a 0.2 μ m filtered solution containing 20
	mM sodium citrate and 0.2 M NaCl, pH 4.5.
Purity	Molecular weight
>98% as determined by SDS-PAGE analysis.	The protein has a calculated MW of 13.75 kDa.
	The protein migrates as 10 kDa under reducing
	condition (SDS-PAGE analysis).
Activity	Endotoxin level
Measure by its ability to induce alkaline phosphatase	<0.1 EU per 1 μ g of the protein by the LAL method.
production by ATDC5 cells. The ED ₅₀ for this effect is	
<2.2 ng/mL.	
Form	

Lyophilized

Background

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2<u>5366</u>77

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw



Background

Bone morphogenetic protein 16 (BMP-16) predicts a molecular mass of 18 kDa. BMPs are multi-functional Growth Factorss that belong to the transforming Growth Factors beta (TGF-β) superfamily. BMPs initiate signaling from the cell surface by binding to two different receptors (R: Type I and II). The heterodimeric formation of type I R and II R may occur before or after BMP binding, inducing signal transduction pathways through SMADs.

Synonyms

Bone Morphogenetic Protein-16, BMP-16

The product is shipped with polar packs. Upon receipt,

store it immediately at -20°C or lower for long term

Sequence Note

His238-Leu347

Shipping

storage.

Uniprot ID

#Q96S42

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in 4 mM HCl to a concentration of 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Image

Tainan Headquarter

Innovation & Research Center

CLD Center


& +886-6-2536677


& +886-6-2536677

☑ bd@leadgene.com.tw

& +886-2-27065528







Human BMP-16, His Tag, E. coli (LDG114PHE) induced alkaline phosphatase production by ATDC5 cells, with the ED50 at 2.119 ng/mL.

SDS-PAGE analysis of recombinant human BMP-16.

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-253<u>6</u>677

& +886-2-27065<u>52</u>8

& +886-6-2536677

☑ bd@leadgene.com.tw