

Human TL1A, His Tag, E. coli

Catalog Number LDG126

Package

LDG126PHE

5 μg / 20 μg / 100 μg / Customized package

For full product information, images and publications, please visit our website.



Specifications

Species of Origin	Expression system
Human	Escherichia coli
Affinity Tag	Buffer
His Tag (C-term)	Lyophilized from a 0.2 μ m filtered solution of PBS, pH
	8.0.
Purity	Molecular weight
>98% as determined by SDS-PAGE analysis.	The protein has a calculated MW of 22.95 kDa.
	The protein migrates as 24 kDa under reducing
	condition (SDS-PAGE analysis).
Activity	Endotoxin level
Activity	
Measure by its ability to induce TF-1 cells proliferation.	<0.1 EU per 1 μ g of the protein by the LAL method.
The ED₅₀ for this effect is <0.2 ng/mL.	
Form	

Lyophilized

Background

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw



Background

TL1A also known as VEGI and TNFSF15, which belongs to TNF ligand superfamily. TL1A is a 28 kDa type II membrane protein containing 251 residues that predominantly expressed in endothelial cells. TL1A is able to promote NFkB activation in cells by interacting with DR3. Besides, TL1A can also induces cell apoptosis through activating caspases. Additionally, the responsiveness of IL-2 from activated T cells has been shown to induce by TL1A treatment.

Synonyms

TNF-Like Ligand 1A, TL1A, Vascular endothelial cell growth inhibitor, Tumor necrosis factor ligand superfamily member 15

The product is shipped with polar packs. Upon receipt,

store it immediately at -20°C or lower for long term

Sequence Note

Gln58-Leu251

Shipping

storage.

Uniprot ID

#O95150

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration of 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 2 weeks under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Image

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-6-2536677

☑ bd@leadgene.com.tw

& +886-2-27065528



kDa	М
	= 1
75	
60	
45	-
35	
25	
15	
10	-

SDS-PAGE analysis of recombinant human TL1A.

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw