

## Human Galectin-3, His Tag, E. coli

**Catalog Number** LDG142PHE

**Package** 5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



### Specifications

#### Species of Origin

Human

#### Affinity Tag

His Tag (N-term)

#### Purity

>95% as determined by SDS-PAGE analysis.

#### Activity

Measured by its ability to chemoattract human PBMC using a concentration range of 2.5-25 µg/mL.

Note: Results may vary from different PBMC donors.

#### Form

Lyophilized

#### Expression system

Escherichia coli

#### Buffer

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

#### Molecular weight

The protein has a calculated MW of 27 kDa. The protein migrates as 32 kDa under reducing condition (SDS-PAGE analysis).

#### Endotoxin level

<0.1 EU per 1 µg of the protein by the LAL method.

### Background

#### Tainan Headquarter

+886-6-2536677 +886-6-2531536

info@leadgene.com.tw

#### Taipei Office

+886-2-27065528

### Background

Galectin-3 (Gal-3) is one of the lectin family members, which is the only chimera-type galectin, containing one carbohydrate recognition domain (CRD) connected to a long, flexible N-terminal domain. The C-terminal CRD is responsible for  $\beta$ -galactoside binding, and the N-terminal domain is essential for its multimerization, and interaction with other intracellular proteins. Galectin-3 is predominantly presented in the cytoplasm and expressed on the cell surface, and then often secreted into biological fluids, such as serum and urine. Numerous studies have indicated that galectin-3 plays a crucial role in cell proliferation, apoptosis, gene expression, immune surveillance, inflammation, fibrosis, and host defense.

### Uniprot ID

#P17931

### Synonyms

Gal-3, 35 kDa lectin, Carbohydrate-binding protein 35, CBP 35, Galactose-specific lectin 3, Galactoside-binding protein, GALBP, IgE-binding protein, L-31, Laminin-binding protein, Lectin L-29, Mac-2 antigen

### Sequence Note

Ala2-Ile250

## Instruction

### Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration not less than 200  $\mu$ g/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

### Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

### Tainan Headquarter

+886-6-2536677 +886-6-2531536

info@leadgene.com.tw

### Taipei Office

+886-2-27065528


### Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

### Image



SDS-PAGE analysis of recombinant human Galectin-3.

**Disclaimer :** For Research Use or Further Manufacturing Only.

#### Tainan Headquarter

+886-6-2536677 +886-6-2531536

info@leadgene.com.tw

#### Taipei Office

+886-2-27065528