

Human CCL4, Tag Free, E. coli

Catalog Number	LDG166PHE
Package	5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Specifications

Species of Origin

Human

Expression system

Escherichia coli

Affinity Tag

Tag Free

Buffer

Lyophilized from a 0.2 µm filtered solution containing 50 mM Tris and 150 mM NaCl, pH 8.5.

Purity

>98% as determined by SDS-PAGE analysis.

Molecular weight

The protein has a calculated MW of 7.75 kDa.
The protein migrates as 11 kDa under reducing condition (SDS-PAGE analysis).

Activity

Measure by its ability to chemoattract human THP1 cells using a concentration range of 10-100 ng/mL.
Activity analysis of recombinant Human CCL4, Tag Free, E. coli.

Endotoxin level

<0.1 EU per 1 µg of the protein by the LAL method.

Form

Lyophilized

Background

Tainan Headquarter

+886-6-2536677

bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677

Background

C-C Motif Chemokine Ligand 4 (CCL4) is a 7.66 kDa cytokine with 69 amino acid residues. CCL4, also named macrophage inflammatory protein-1 β (MIP-1 β), is mainly secreted from neutrophils, monocytes, B cells, T cells, fibroblasts, endothelial cells and epithelial cells. CCL4 recruits various immune cells like natural killer cells, monocytes, and neutrophils when CCL4 binds to CCR5. In addition, CCL4 mediates lymphocyte adhesion by Rac1 /Cdc42 signaling pathway and plays critical roles in different biological functions like maintenance of cell polarity, regulation of calcium ion transport and response to viruses.

Uniprot ID

#P13236

Synonyms

Lymphocyte activation gene 1 protein, LAG-1, Macrophage inflammatory protein 1-beta, MIP-1-beta, Monocyte adherence-induced protein 5-alpha, Small-inducible cytokine A4-like, C-C motif chemokine 4-like

Sequence Note

Ala24-Asn92

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration of 200 μ g/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Image

Tainan Headquarter

+886-6-2536677


bd@leadgene.com.tw

Innovation & Research Center


+886-2-27065528

CLD Center

+886-6-2536677



Measure by its ability to chemoattract human THP1 cells using a concentration range of 10-100 ng/mL. Activity analysis of recombinant Human CCL4, Tag Free, E. coli.



SDS-PAGE analysis of recombinant human CCL4.

Disclaimer : For Research Use or Further Manufacturing Only.